92.6%,此项结果提示病理科医师可以根据TTF-1或者SPA的免疫组化结果对EGFR突变状态进行简单的预测,同样临床医师也可初步判断患者的预后以及化疗效果。

以上美国、韩国、法国以及我们的数据,均来自肺腺癌的研究,在肺癌其他类型,如鳞癌、大细胞等非小细胞肺癌中TTF-1表达与EGFR突变状态的关系并不明确,需要更进一步的研究。

参考文献:

[1] Pao W, Miller V, Zakowski M, *et al*. EGF receptor gene mutations are common in lung cancers from "never smokers" and are associated with sensitivity of tumors to gefitinib and erlotinib[J]. Proc Natl Acad Sci U S A, 2004, 101(36): 13306-13311.

[2] Eberhard D A, Johnson B E, Amler L C, *et al*. Mutations in the epidermal growth factor receptor and in KRAS are predictive and prognostic indicators in patients with non-small-cell lung cancer treated with chemotherapy alone and in combination with erlotinib[J]. J Clin Oncol, 2005, 23(25): 5900-5909.

[3] Tsao M S, Sakurada A, Ding K, *et al*. Prognostic and predictive value of epidermal growth factor receptor tyrosine kinase domain mutation status and gene copy number for adjuvant chemotherapy in non-small cell lung cancer[J]. J Thorac Oncol, 2011, 6(1): 139-147.

[4] 奉水东, 谭红专. 肺癌EGFR基因突变检测方法的研究进展[J]. 中国肺癌杂志, 2008, 11(3): 462-464.

[5] Fabbro D, Di-Loreto C, Stamerra O, *et al*. TTF-1 gene expression in human lung tumours[J]. Eur J Cancer, 1996, 32A(3): 512-517.

[6] Boggaram V. Thyroid transcription factor-1 (TTF-1/Nkx2.1/TITF1) gene regulation in the lung[J]. Clin Sci (Lond), 2009, 116(1): 27-35.

[7] Kaufmann O, Dietel M. Thyroid transcription factor-1 is the superior immunohistochemical marker for pulmonary adenocarcinomas and large cell carcinomas compared to surfactant proteins A and B[J]. Histopathology, 2000, 36(1): 8-16.

[8] Tang X, Kadara H, Behrens C, *et al*. Abnormalities of the TITF-1 lineage-specific oncogene in NSCLC: implications in lung cancer pathogenesis and prognosis[J]. Clin Cancer Res, 2011, 17(8): 2434-2443.

[9] Chang Y L, Lee Y C, Liao W Y, *et al*. The utility and limitation of thyroid transcription factor-1 protein in primary and metastatic pulmonary neoplasms[J]. Lung Cancer, 2004, 44(2): 149-157.

[10] Berghmans T, Paesmans M, Mascaux C, *et al*. Thyroid transcription factor 1--a new prognostic factor in lung cancer: a meta-analysis[J]. Ann Oncol, 2006, 17(11): 1673-1676.

[11] Somaiah N, Garrett-Mayer E, Huang X, *et al*. Use of negative thyroid transcription factor (TTF-1) status to predict for negative epidermal growth factor receptor (EGFR) mutations (Mts) status with a high negative predictive value (NPV) in patients (pts) with adenocarcinomas (AC) of the lung[J]. J Clin Oncol, 2011, 29(Suppl): 7530.

[12] Sun P L, Seol H, Lee H J, *et al*. High incidence of EGFR mutations in Korean men smokers with no intratumoral heterogeneity of lung adenocarcinomas: correlation with histologic subtypes, EGFR/TTF-1 expressions, and clinical features[J]. J Thorac Oncol, 2012, 7(2): 323-330.

(**收稿**:2012-07-05;**修回**:2012-09-16)

(编辑 汪勤俭)

"神经创伤救治的转化医学研究"重点号2012年第23期即将出版

专家述评

1 神经创伤救治的转化医学思考…… 冯 华

论著

1 去铁胺对大鼠大脑冲击伤的治疗作用…… 张礼均等
2 瞬时受体电位通道6在大鼠大脑冲击伤后铁代谢中的作用及机制研究…… 张礼均等
3 蜕皮甾酮对大鼠脑组织自由基损伤的保护作用…… 高诗豪等
4 草麻黄中补体抑制单体防治大鼠脑冲击伤疗效的研究…… 魏林节等

本刊编辑部