
维普资讯 http://www.cqvip.com





a:支架置入术前; b:支架置入术后

**图1 支架置入术前、后主动脉造影结果**

更大直径的球囊扩张使支架内径增加以解除缩窄。因此,对于主动脉缩窄合并动脉导管未闭的患者进行覆膜支架置入治疗时,只要指征选择恰当,完全可以达到有效根治之目的。

[参考文献]

[1] 朱晓东. 心脏外科基础图谱[M]. 第2版. 北京:人民卫生出版社,2002.444-445.

[2] 刘玉清. 心血管病影像诊断学[M]. 第2版. 合肥:安徽科学技术出版社,2000.577.

[3] 刘 芳,黄国英,梁雪村,等. 主动脉缩窄96例临床分析[J]. 中华医学杂志,2006,86(26):1854-1856.

[4] Rao PS, Chopra PS. Role of balloon angioplasty in the treatment of aortic coarctation[J]. Ann Thorac Surg,1991,52(3):621-631.

[5] Corno AF, Botta U, Hurni M, et al. Surgery for aortic coarctation: a 30 years experience[J]. Eur J Cardiothorac Surg,2001,20(2):1202-1206.

[6] Dietl CA, Torres AR, Favaloro RG, et al. Right submammarian thoracotomy in female patients with atrial septal defects and anomalous pulmonary venous connections. Comparison between the transpectoral and subpectoral approaches[J]. J Thorac Cardiovasc Surg,1992,104(3):723-737.

[7] Mullins CE, O'Laughlin MP, Vick GW, et al. Implantation of balloon-expandable intravascular grafts by catheterization in pulmonary arteries and systemic veins[J]. Circulation,1988,77(1):188-199.

[8] O'Laughlin MP, Perry SB, Lock JE, et al. Use of endovascular stents in congenital heart disease[J]. Circulation, 1991, 83(6):1923-1939.

[9] Veith FJ, Abbott WM, Yao JS, et al. Guidelines for development and use of transluminally placed endovascular prosthetic grafts in the arterial system. Endovascular Graft Committee[J]. J Vasc Interv Radiol, 1995,6(3):477-491.

[10] Chuter TA,Green RM,Ouriel K,et al. Transfemoral endovascular aortic graft placement[J]. J Vasc Surg,1993,18(2):185-197.

[11] Ewert P, Schubert S, Peters B, et al. The CP stent-short, long, covered-for the treatment of aortic cortic cortictatuon, stenosis of pulmonary arteries and caval veins, and Fontan anastomosis in children and adults: an evaluation of 60 stents in 53 patients[J]. Heart, 2005,91(7):948-953.

[12] 舒茂琴,宋治远,刘建平,等. 主动脉缩窄合并动脉导管未闭的介入治疗(附1例报道)[J]. 重庆医学,2006,35(4):300-301.

[13] 俞飞成,黄连军,吴文辉,等. 带膜支架置入治疗主动脉缩窄合并动脉导管未闭一例[J]. 中华心血管病杂志,2006,34(6):563.

[14] Pedra CAC,Fontes VF,Esteves CA,et al. Use of covered stents in the management of coarctation of the aorta[J]. Pediatric Cardiology, 2005,26(4):431-439.

(编辑:郭光金)

## 关于单位符号的使用

各种单位除在无数值的叙述性文字中和科普读物中可使用中文名称或中文符号外,均应使用单位符合。单位符号都是正体拉丁和希腊字母,无复数形式,符号后不附加圆点(正常语法名字尾标点符号除外)。来源于人名的单位,例如:Pa、Gy、W等和$10^6$以上的词头,例如:M、G、T等的首字母大写以外均为小写,"升"的符号可用大写(L)也可用小写(l),一般在单位使用不加词头时用大写。数值与单位符号之间、单位符号与相邻的其他符号(例如:化学元素符号)之间应留半个阿拉伯数字空隙,词头与其所组合的单位符号之间不需空隙。词头不能单独使用,也不能重叠使用(例如:"mμm"应为"nm")。

(本刊编辑部)