证医学研究等还有待进一步发展与完善。

参考文献：

[1] Lourenco-Matharu L, Roberts G J. Oral sedation for dental treatment in young children in a hospital setting[J]. Br Dent J, 2010, 209(7): E12.

[2] Garip H, Satilmis T, Dergin G, *et al*. Effects of Midazolam/Low Dose Ketamine Conscious Intravenous Sedation on Pain, Swelling, and Trismus After Surgical Extraction of Third Molars[J]. J Oral and Maxillofac Surg, 2011 Jan 6. [Epub ahead of print]

[3] Danda A K, Krishna-Tatiparthi M, Narayanan V, *et al*. Influence of primary and secondary closure of surgical wound after impacted mandibular third molar removal on postoperative pain and swelling-a comparative and split mouth study[J]. J Oral Maxillofac Surg, 2010, 68(2): 309 - 312.

[4] Ryding H A, Murphy H J. Use of nitrous oxide and oxygen for conscious sedation to manage pain and anxiety[J]. J Can Dent Assoc, 2007, 73(8): 711.

[5] Emmanouil D E, Quock R M. Advances in understanding the actions of nitrous oxide[J]. Anesth Prog, 2007, 54(1): 9 - 18.

[6] Emmanouil D E, Dickens A S, Heckert R W, *et al*. Nitrous oxide-antinociception is mediated by opioid receptors and nitric oxide in the periaqueductal gray region of the midbrain[J]. Eur Neuropsychopharmacol, 2008, 18(3): 194 - 199.

[7] Fujinaga M, Maze M. Neurobiology of nitrous oxide-induced antinociceptive effects[J]. Mol Neurobiol, 2002, 25(2): 167 - 189.

[8] Georgiev S K, Kohno T, Ikoma M, *et al*. Nitrous oxide inhibits glutamatergic transmission in spinal dorsal horn neurons[J]. Pain, 2008, 134(1/2): 24 - 31.

[9] Gruss M, Bushell T J, Bright D P, *et al*. Two-pore-domain K^+ channels are a novel target for the anesthetic gases xenon, nitrous oxide, and cyclopropane[J]. Mol Pharmacol, 2004, 65(2): 443 - 452.

[10] Farrell M K, Drake G J, Rucker D, *et al*. Creation of a registered nurse-administered nitrous oxide sedation program for radiology and beyond[J]. Pediatr Nurs, 2008, 34(1): 29 - 36.

[11] Baygin O, Bodur H, Isik B. Effectiveness of premedication agents administered prior to nitrous oxide/oxygen[J]. Eur J Anaesthesiol, 2010, 27(4): 341 - 346.

[12] Wood M. The safety and efficacy of intranasal midazolam sedation combined with inhalation sedation with nitrous oxide and oxygen in paediatric dental patients as an alternative to general anaesthesia[J]. SAAD Dig, 2010, 26: 12 - 22.

[13] Okushima K, Kohjitani A, Asano Y, *et al*. Inhalational conscious sedation with nitrous oxide enhances the cardiac parasympathetic component of heart rate variability[J]. Oral Surg Oral Med Oral Pathol Oral Radiol Endod, 2008, 106(6): e1 - e5.

[14] Scheffer A C, Schuurmans M J, vanDijk N, *et al*. Reliablity and validity of the visual analogue scale for fear of falling in older persons[J]. J Am Geriatr Soc, 2010, 58(11): 2228 - 2230.

[15] 熊章荣，唐延先. 多瑞吉联合罗哌卡因用于腹腔镜胆囊切除术后镇痛的临床研究[J]. 第三军医大学学报，2007，29(7)：643 - 645.

[16] Goerig M, Schulte-am-Esch J. Early contributions for the development of nitrous oxide-oxygen anesthesia in central Europe[J]. Anaesthesiol Reanim, 2002, 27(2): 42 - 53.

[17] Cote C J, Wilson S. Guidelines for monitoring and management of pediatric patients during and after sedation for diagnostic and therapeutic procedures: an update[J]. Pediatrics, 2006, 118(6): 2587 - 2602.

（收稿：2010-12-20 修回：2011-02-09）

（编辑 龙 亮）

2011国际暨全国第十一届头颈肿瘤学术大会征文通知

由中国抗癌协会头颈肿瘤专业委员会主办的2011国际暨全国第十一届头颈肿瘤学术大会定于2011年9月16 - 18日在杭州召开。本届大会已被确定为国家级继续教育项目，参会者将获得继续教育学分（Ⅰ类）10分。

征文内容：头颈肿瘤的综合序列治疗、头颈肿瘤的微创治疗、头颈肿瘤（包括耳鼻咽喉、口腔等）临床与基础研究、诊治经验，新技术、新方法介绍，护理及预防、康复等方面论著、个案等。

来稿要求：①论文须未公开发表过，正文5000字以内；②500字左右中英文摘要（包括目的、方法、结果、结论）及关键词；③正文与摘要分开、另纸打印；④注明第一作者姓名、单位、地址、邮编，并自留底稿。

提交方式：①以电子邮件的方式发至headneck2011@163.com，请注明“2011国际暨全国第十一届头颈肿瘤学术大会征文”！联系人：赏金标、陈超、郑传铭，联系方式：赏金标13605807802、陈超13958013018、郑传铭13868006966。②邮寄至杭州市武林广场东侧省科协大楼10楼1005室（邮编：310003），加盖单位公章。③截稿日期：2011年6月30日，以电子邮件或邮寄时间为准。

大会组织委员会和学术委员会将邀请、组织有关专家认真审读投稿论文，评选出优秀论文给予表彰和奖励，并作大会报告或以壁报形式在会议期间展览交流，部分论文将向SCI收录的期刊推荐。